

The Role of the Hippocampus in Learning and Memory

Yerui Wang

Department of Biological Science and Technology, Tokyo University of Science, Tokyo, 162-8601, Japan

Corresponding author: 8325011@ed.tus.ac.jp

Abstract

Memory and learning represent fundamental operations of the cerebral cortex, with the hippocampus playing a pivotal role in these processes. Synaptic plasticity is currently recognized as a primary mechanism through which neurons acquire and retain information. Recently, numerous investigations have broadened our comprehension regarding hippocampal synaptic plasticity. Nevertheless, the integrated functions of diverse hippocampal sectors and various types of synaptic modifications remain largely unexplored. This article provides an overview of contemporary research concerning hippocampal synaptic plasticity in cognition. Initially, it describes the primary architecture and roles of the hippocampus while clarifying how distinct hippocampal segments contribute to memory consolidation. Subsequently, it delineates the general concepts of synaptic plasticity before concentrating on the molecular underpinnings of long-term potentiation (LTP) and long-term depression (LTD). This manuscript will also discuss the synergy between these two forms of plasticity in modulating neuronal circuits and facilitating mnemonic processes. In total, it is evident that learning and memory emerge from synchronized activities across multiple hippocampal components and equilibrium shifts in synaptic states. This research elucidates how synaptic modifications in the hippocampus link to cognitive functions. It serves as the foundation for subsequent inquiries into cerebral memory impairments. Prospective investigations will examine the associations among all hippocampal sub-fields and analyze the molecular mechanisms of synaptic plasticity within normal and pathological cerebral states.

Key words

Hippocampus; LTP(long-term potentiation); LTD(long-term depression)

1. Introduction

Acquisition and retention represent fundamental

roles of the neural cortex. These processes assist individuals in grasping novel information, recalling positive experiences, and utilizing previous insights within their

everyday routines. Numerous cerebral regions participate in this procedure, yet the hippocampus plays a crucial part. It aids in forming fresh recollections while also accounting for the creation of spatial and episodic cognitions [1, 2].

The Hippocampus possesses no singular role. Rather, it comprises several distinct sectors, including the dentate gyrus (DG), CA1 and CA3. Each of these areas possesses unique architectures and executes diverse tasks during the processes of learning and memory. Collectively, they manage, conserve and recall data [2,3]. Numerous investigations involving both humans and animals have demonstrated that impairment of the hippocampus might result in severe cognitive deficits. Consequently, the hippocampus has started capturing the focus of neuroscientists during the past few decades [2,3].

Studies indicate that the hippocampus assists in cognitive learning and memory. Within these diverse mechanisms, synaptic plasticity is viewed as a highly significant factor [1,3,4]. The process of altering the intensity of neuronal linkages after activation is termed synaptic plasticity. It enables experiential learning and preserves novel insights within the neural tissue. Two primary forms of long-term synaptic plasticity include long-term potentiation (LTP) and long-term depression (LTD). LTP enhances the interaction between two neurons, while LTD diminishes it. Both are essential for reinforcing crucial data in the brain and discarding unnecessary synaptic pathways [1,4,5].

Dopamine represents an additional neuromodulator capable of modifying synaptic efficacy, thus impacting both cognition and recollection. Lately, various investigations have revealed that regulatory patterns across distinct hippocampal sectors differ. The DG, CA3 and CA1 compartments perform unique roles in synaptic flexibility and mnemonic processes within separate zones [3].

Recently, significant progress has emerged in investigations regarding synaptic plasticity within the hippocampus. Utilizing advanced techniques, investigators have acquired deeper insights into how neural cells interact and establish memories. Concurrently, numerous inquiries have likewise demonstrated that distorted synaptic plasticity is linked to diverse neurodegenerative conditions, including Alzheimer's disease (AD), Parkinson's disease (PD) and senile cognitive impairment.

Consequently, exploring hippocampal synaptic plasticity may assist us in comprehending brain mechanisms more effectively and creating novel therapeutic strategies for cognitive disorders [3,6].

This article aims to investigate the role of synaptic plasticity in the hippocampus regarding learning and memory. Initially, it will present the fundamental architecture and functions of the hippocampus. Subsequently, it will cover the general mechanisms of synaptic plasticity and specifically emphasize LTP and LTD. Ultimately, there is a review of the research findings and future directions. The preceding discussion shall briefly outline the current understanding of hippocampal synaptic plasticity and clarify why it is essential for learning and memory.

2. The Hippocampus and Memory Formation

2.1 Structure of the Hippocampus

Located within the medial temporal lobe, the hippocampus is a primary brain region specifically tasked with acquiring and retaining knowledge [5]. Despite being commonly regarded as a unified entity, it is composed of distinct sectors performing unique functions. These primary segments include DG, CA3, CA1, and the subiculum [2,5]. These sub-regions are interconnected via a highly structured neural network. Instead of operating independently, they collectively collaborate to manage, convey, and preserve data.

The sub-regions exhibit distinct characteristics regarding their memory capacity. Dentate cells facilitate the organization of analogous experiences within the brain into distinct patterns and differentiate between them [2]. Consequently, there will be less overlap in stored memories. The CA3 area is linked to the dentate gyrus, and through pattern completion, it can reconstruct complete recollections from fragmented data [2]. The CA1 area gathers information from CA3 and various brain segments and subsequently transmits the refined data to the cortex via the subiculum [5]. Since every subregion serves a unique function, injury in diverse locations could lead to varied categories of amnesia. Thus, to understand the mechanisms of learning and memory,

it is essential to comprehend the architecture of the hippocampus.

2.2 Hippocampus-dependent Learning and Memory

The hippocampus is essential for learning and memory processes that depend on its operation. Nevertheless, it primarily participates in the creation of fresh memories rather than being involved in the enduring retention of every recollection [5]. Among various forms of memory, spatial and episodic memory are the most intimately associated with hippocampal activity [2]. Spatial memory assists individuals and creatures in identifying their positions and navigating through particular environments; episodic memory is utilized to retrieve individual encounters and specific occurrences. These two categories of memory have been extensively employed to investigate the role of the hippocampus in both humans and animals [2, 5].

Numerous researches have demonstrated that the hippocampus participates in the mechanisms of learning and recollection. A prominent example involves patient H.M., who underwent a procedure removing portions of his medial temporal cortex and hippocampus; he subsequently lost the ability to create novel long-term recollections [5]. Nevertheless, his transient memory and numerous pre-operative memories remained largely functional [5]. Consequently, it was suggested that the hippocampus preserves and consolidates new long-term recollections instead of merely holding them temporarily. This further illustrated that the neural processes of transient and permanent memory are distinct [5].

Numerous investigations have likewise been performed upon mammals. Rodents are capable of learning to locate a concealed platform within the Morris Water Maze through spatial cues. Subjects possessing hippocampal damage require greater durations to reach the target and exhibit diminished spatial cognition relative to normal subjects [5]. Consistent findings were derived from the radial arm maze test; specifically, hippocampal deficits result in increased navigational mistakes and deteriorated recollection [5]. Collectively, these previous inquiries indicate that an intact hippocampus is essential for cognition and memory.

Recently, various researches have revealed that

memory establishment is not confined to a single site within the hippocampus. Rather, it represents the cumulative impact of numerous regions throughout this structure [2]. The DG, CA3 plus CA1 possess distinct yet interconnected functions in cognition. Collectively, they facilitate the encoding, stabilization and retrieval of recollections in the hippocampus [2]. Consequently, hippocampus-reliant learning and memory must be viewed as the outcome of synergy among multiple subfields of the hippocampus, instead of the responsibility of a specific location. To understand how the hippocampal network is modified during training, Research must examine synaptic plasticity more rigorously in the subsequent paragraphs.

3. Synaptic Plasticity in the Hippocampus

3.1 Basic Concepts of Synaptic Plasticity

The hippocampus facilitates learning and recall, and this function is executed through modifications in neurons. A primary vital alteration involves synaptic plasticity. Synaptic plasticity represents the modulation in synaptic strength resulting from neuronal firing at different intervals [1,4]. This property can be adjusted via experience, and it is typically regarded as one of the fundamental building blocks of learning and memory at the cellular scale [1].

Synaptic Plasticity is a continuous process rather than a binary state. It fluctuates constantly as the neural network acquires fresh life experiences. The two primary categories of synaptic modification are LTP and LTD [1,4]. While both involve adjustments to synaptic potency, they exert contrasting influences on the efficiency of neural signaling. Following multiple high-frequency pulses, LTP enhances the power of synaptic coupling, whereas LTD diminishes it through sustained low-frequency firing or diverse patterns of neuronal engagement [1, 4]. These mechanisms are essential for cognition and recollection because the cerebral cortex must solidify the links of beneficial neurons while diminishing or discarding those that remain unnecessary [1].

The occurrence of LTP and LTD is primarily attributed to fluctuations in cytosolic calcium (Ca^{2+}) levels

and the engagement of numerous signaling cascades [1,4]. An elevation in intracellular calcium follows intense synaptic firing, which subsequently triggers the mechanism for synaptic potentiation. Conversely, feeble or extended stimuli merely cause a minor, transient spike in Ca^{2+} without activating the specific signaling pathway for synaptic enhancement [1, 4]. Modifications in molecules at the molecular level offer the basis for enduring shifts in synaptic performance resulting from cognitive processes. The specific molecular mechanisms underlying LTP and LTD are discussed in the subsequent sections.

Numerous factors trigger synaptic plasticity. Specifically, several are neurotransmitters governing neural dynamics. Dopamine modulates synaptic plasticity within the hippocampus, influencing memory consolidation across distinct hippocampal zones [3]. Lately, certain research has discovered that such modulation varies across different hippocampal subfields. The DG, CA3, and CA1 exhibit diverse synaptic plasticity profiles and play distinct roles in mnemonic processing [2,3]. Consequently, it is inferred that synaptic plasticity operates at every tier and numerous synapses. Instead, this represents an adaptation arising from the coordinated function of various hippocampal pathways.

Synaptic modifications offer the fundamental cellular framework for cognition and recollection, thus remaining a major subject of inquiry within neurobiology currently [1]. The subsequent paragraphs will outline the molecular pathways of LTP and LTD and clarify how these mechanisms facilitate hippocampal-reliant memory and learning.

3.2 LTP

LTP represents a primary category of synaptic modulation. Initially identified within the hippocampus, this phenomenon serves as a prototypical model for various cognitive dysfunctions [1,4]. LTP denotes an enduring strengthening of neural connections triggered by numerous occurrences of high-frequency input [1]. Since it leads to persistent modifications in neuronal signaling patterns, numerous scientists have postulated that LTP constitutes a fundamental cellular basis for the establishment of memories [1,5].

LTP initiates following the discharge of glutamate from presynaptic cells into the synaptic cleft following

intense-frequency activation [6]. Glutamate initially attaches to AMPA-type receptors located on the postsynaptic surface. Consequently, sodium ions (Na^+) may penetrate the postsynaptic cell and trigger membrane depolarization [6]. This depolarization subsequently removes the magnesium (Mg^{2+}) inhibition from the NMDA receptor. Thus, NMDA receptors become activated and calcium ions (Ca^{2+}) flow into the postsynaptic cell [6,7]. The initial phase of LTP formation involves an elevated concentration of intracellular Ca^{2+} .

Rising intracellular calcium levels trigger numerous signaling pathways within the postsynaptic neuron [8, 9]. Specifically, among these, calcium/calmodulin-dependent protein kinase II (CaMKII) serves as a primary element in LTP [6]. When activated, CaMKII enhances both the quantity and function of AMPA receptors located on the postsynaptic membrane [6,8]. An elevated density of AMPA receptors bolsters the postsynaptic neuron's reaction to glutamate subsequently. Consequently, the strength and velocity of synaptic conveyance are elevated [6]. This process is quite enduring, and as a cellular entity, it possesses the capacity to retain data for an extended duration [1,6].

Molecular modifications within cerebral neural networks occur simultaneously with LTP. Enhanced synapses exhibit superior connectivity among neurons. Consequently, information transfer becomes more efficient within the hippocampal architecture [5]. LTP patterns vary across distinct hippocampal regions as well. DG, CA3, and CA1 display diverse synaptic plasticity profiles and play roles in memory formation differently [2,3]. Furthermore, dopamine modulates LTP in the CA1 region via dopaminergic pathways, and such alterations facilitate learning and memory for various types of behavior [3].

Numerous researches have revealed that standard LTP is linked to effective learning and recollection. LTP deficiency typically causes a decrease in cognitive ability. Dysfunctional LTP has likewise been identified in various neurodegenerative conditions, including Alzheimer's disease and senile cognitive impairment [5]. Consequently, investigating the molecular basis of LTP is essential to comprehend how memories are established and where to locate novel therapeutics for deficits of memory.

LTP enhances synaptic links and facilitates mem-

ory establishment, yet it does not serve as the exclusive foundation for healthy brain activity. Decrease the density of redundant synapses within the brain and maintain the equilibrium of neuronal circuits. The primary technique to accomplish this involves LTD; it shall be discussed in the subsequent paragraph.

3.3 LTD

LTD represents another category of synaptic plasticity. This process involves a sustained reduction in synaptic efficacy, whereas LTP signifies an enhancement [1,4]. Despite these mechanisms exhibiting contrasting impacts, both are essential for the proper operation of the cerebral cortex. LTP consolidates robust neuronal links, while LTD diminishes fragile or non-utilized ones. Collectively, they enable perpetual modification of neural networks within the brain during the acquisition and retention of knowledge [10].

LTD typically denotes the consequence of extended low-frequency activation or specific neural firing sequences [6]. Glutamate consequently continues to engage AMPA and NMDA ion channels. Nevertheless, the magnitude of postsynaptic depolarization remains below levels observed in LTP [6]. Consequently, NMDA receptors trigger a minor yet persistent elevation in cytosolic calcium (Ca^{2+}). Such a minimal calcium influx then activates phosphatases rather than kinases [6,9]. The aforementioned signaling cascades diminish the density of AMPA receptors at the synaptic junction, thereby lowering synaptic efficacy [6,10]. LTD does not signify the termination of synaptic communication. It merely decreases the strength of synaptic signaling and modifies the overall pattern of neuronal firing.

LTD is essential for maintaining brain function in a healthy condition. Should synaptic associations remain perpetually potent, neural pathways might become overwhelmed and lose their capacity to acquire fresh data [10]. LTD may diminish the intensity of certain feeble synapses to regulate the general degree of synaptic engagement. Adjust Synapses to Enhance the Flexibility of the Neural Framework. Consequently, the cortex can eliminate redundant data and preserve vital recollections [10]. Thus, typical learning involves not only fortifying synaptic bonds but also precisely diminishing specific ones at specific intervals.

LTD has been identified in various regions of the

hippocampus recently [2,3]. The DG, CA3, and CA1 exhibit distinct profiles of synaptic plasticity and react to neuronal firing in diverse manners [2]. Dopamine also modulates LTD within the hippocampus through dopaminergic signaling pathways, although the magnitude of such influence fluctuates depending on the specific behavioral context [3]. Consequently, it is inferred that LTD represents a controlled mechanism rather than a simple reduction in synaptic efficacy.

Dysregulated LTD is linked to numerous neurodegenerative conditions, including Alzheimer's disease and senile cognitive deterioration [5]. Shifts in the LTP/LTD equilibrium may hinder memory consolidation and disrupt the healthy operation of neural circuits. Consequently, to preserve cerebral wellness, these two forms of synaptic flexibility must remain in equilibrium. LTP and LTD function synergistically to establish the molecular foundation for cognition and adjustment by modifying synaptic linkages within the brain according to environmental stimuli.

4. Conclusion

The preceding enumeration encompasses the manifestations of hippocampal synaptic plasticity within learning and memory. Initially, it presented the fundamental architecture and operations of the hippocampus while clarifying the roles of diverse hippocampal regions in memory processes. Subsequently, it summarized the fundamental concepts of synaptic plasticity and further examined the molecular mechanisms underlying LTP and LTD. Collectively, these findings demonstrate that synaptic plasticity serves as the cellular basis for learning and memory, enabling the brain to modify itself in response to novel experiences.

This research demonstrates that acquisition and recollection are far more intricate than previously believed. These processes are not confined to a specific region within the hippocampus nor triggered by a unique chemical. Instead, they are regulated by diverse components of the hippocampus and multiple types of synaptic adaptability. LTP and LTD do not function in isolation. Combined, they manage the equilibrium of neural circuits and facilitate the encoding, stabilization, and refinement of remembrance. Jointly, they enable us to understand how to establish and retain cognition over

duration. It assists us in investigating memory-associated neurological disorders more effectively and creating novel therapies hereafter. Despite certain advancements in recent times, the precise mechanisms of hippocampal synaptic flexibility in the human cerebrum remain incompletely understood. Novel techniques will be utilized subsequently to examine molecules and neurological maladies to enhance care for memory impairments.

References

- [1] Bliss, T. V., & Collingridge, G. L. (1993). Synaptic basis of memory: Long-term potentiation in the hippocampus. *Nature*, 361(6407), 31–39. <https://doi.org/10.1038/361031a0>
- [2] Danieli, K., Guyon, A., & Bethus, I. (2023). Episodic memory development: An overview of diverse hippocampal input trajectories. *Progress in Neuro-Psychopharmacology & Biological Psychiatry*, 126, 110757. <https://doi.org/10.1016/j.pnpbp.2023.110757>
- [3] Tsetsenis, T., Broussard, J. I., & Dani, J. A. (2023). Dopamine control of hippocampal plasticity, learning and memory. *Frontiers in Behavioral Neuroscience*, 16, 1092420. <https://doi.org/10.3389/fnbeh.2022.1092420>
- [4] Crystal, J. D. (2013). Prospective memory. *Current Biology*, 23(17), R750–R751. <https://doi.org/10.1016/j.cub.2013.07.081>
- [5] Kandel, E. R., Koester, J. D., Mack, S. H., & Siegelbaum, S. A. (2021). *Principles of neural science* (6th ed.). McGraw Hill.
- [6] Chen, Q. Y., Li, X. H., & Zhuo, M. (2021). NMDA receptors and synaptic plasticity within the anterior cingulate cortex. *Neuropharmacology*, 197, 108749. <https://doi.org/10.1016/j.neuropharm.2021.108749>
- [7] Collingridge, G. L. (2025). Long-term potentiation in the hippocampus: From magnesium to memory. *Neuroscience*, 578, 126–131. <https://doi.org/10.1016/j.neuroscience.2024.11.069>
- [8] Quintanilla, J., Jia, Y., Pruess, B. S., et al. (2024). Pre- and post-synaptic manifestations of LTP within two branches of one hippocampal input. *Journal of Neuroscience*, 44(10), e1449232024. <https://doi.org/10.1523/JNEUROSCI.1449-23.2024>
- [9] Nicoll, R. A., & Schulman, H. (2023). Synaptic memory and CaMKII. *Physiological Reviews*, 103(4), 2877–2925. <https://doi.org/10.1152/physrev.00034.2022>
- [10] Azarfarin, M., Ghadiri, T., Dadkhah, M., & Sahab-Negah, S. (2024). The interface between cannabinoids and enduring synaptic plasticity: An overview of cognitive encoding and underlying pathways. *Cell Biochemistry and Function*, 42(6), e4100. <https://doi.org/10.1002/cbf.4100>